

# HSDL1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56313

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q3SXM5</a>
<b>Reactivity</b>	Rat, Pig, Dog, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	37002
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human HSDL1
<b>Epitope Specificity</b>	161-260/330
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Mitochondrion.
<b>SIMILARITY</b>	Belongs to the short-chain dehydrogenases/reductases (SDR) family. 17-beta-HSD 3 subfamily.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	<p>HSDL1 is a 330 amino acid protein that belongs to the short-chain dehydrogenases/reductases (SDR) family and 17-beta-HSD 3 subfamily. Localizing to the mitochondrion, HSDL1 is highly expressed in testis and ovary, with lower levels of expression found in thyroid, spinal cord, adrenal gland, heart, placenta, skeletal muscle, small intestine, colon, spleen, prostate and pancreas. HSDL1 interacts with DUSP24 and is encoded by a gene that maps to human chromosome 16q23.3 and mouse chromosome 8 E1. Chromosome 16 encodes over 900 genes in approximately 90 million base pairs, makes up nearly 3% of human cellular DNA, and is associated with a variety of genetic disorders. The rare disorder Rubinstein-Taybi syndrome is associated with chromosome 16 through the CREBBP gene, which encodes a critical CREB binding protein. Crohn's disease is a gastrointestinal inflammatory condition associated with chromosome 16 through the NOD2 gene.</p>

## Additional Information

---

<b>Gene ID</b>	83693
<b>Other Names</b>	Inactive hydroxysteroid dehydrogenase-like protein 1, Short chain dehydrogenase/reductase family 12C member 3, HSDL1, SDR12C3
<b>Target/Specificity</b>	Highly expressed in testis and ovary. Also detected in thyroid, spinal cord, adrenal gland, heart, placenta, skeletal muscle, small intestine, colon, spleen,

prostate and pancreas.

<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	HSDL1
<b>Synonyms</b>	SDR12C3
<b>Cellular Location</b>	Mitochondrion.
<b>Tissue Location</b>	Highly expressed in testis and ovary. Also detected in thyroid, spinal cord, adrenal gland, heart, placenta, skeletal muscle, small intestine, colon, spleen, prostate and pancreas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.