

TRAPPCL2 Rabbit pAb

TRAPPCL2 Rabbit pAb

Catalog # AP56328

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UL33
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16146
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRAPPCL2
Epitope Specificity	21-80/140
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasmic, Endoplasmic reticulum and Golgi Apparatus
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	TRAPPC2L may play a role in vesicular transport from endoplasmic reticulum to Golgi.

Additional Information

Gene ID	51693
Other Names	Trafficking protein particle complex subunit 2-like protein, TRAPPC2L
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TRAPPC2L
Function	Plays a role in vesicular transport from endoplasmic reticulum to Golgi.
Cellular Location	Cytoplasm, perinuclear region. Endoplasmic reticulum. Golgi apparatus


Tissue Location

Expressed in testis, liver, bladder, lung, spleen and brain, several cell lines and primary chondrocytes cell line

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRAPPCL2) Polyclonal Antibody, Unconjugated (AP56328) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.