

INO80 Rabbit pAb

INO80 Rabbit pAb
Catalog # AP56350

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9ULG1
Reactivity	Mouse
Predicted	Rat, Dog, Human, Rabbit, Monkey, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	176753
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human INO80
Epitope Specificity	101-200/1556
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Colocalizes with PCNA at replication forks during S-phase. Recruited to DNA damage sites in a ACTR8-dependent manner.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a subunit of the chromatin remodeling complex, which is classified into subfamilies depending on sequence features apart from the conserved ATPase domain. This protein is the catalytic ATPase subunit of the INO80 chromatin remodeling complex, which is characterized by a DNA-binding domain. This protein is proposed to bind DNA and be recruited by the YY1 transcription factor to activate certain genes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013]

Additional Information

Gene ID	54617
Other Names	Chromatin-remodeling ATPase INO80, hINO80, 3.6.4.-, DNA helicase-related INO80 complex homolog 1, DNA helicase-related protein INO80, INO80 complex subunit A, INO80, INO80A, INOC1, KIAA1259
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	INO80
Synonyms	INO80A, INOC1, KIAA1259
Function	ATPase component of the chromatin remodeling INO80 complex which is involved in transcriptional regulation, DNA replication and DNA repair (PubMed: 16230350 , PubMed: 16298340 , PubMed: 17721549 , PubMed: 20237820 , PubMed: 20855601). Binds DNA (PubMed: 16298340 , PubMed: 21303910). As part of the INO80 complex, remodels chromatin by shifting nucleosomes (PubMed: 16230350 , PubMed: 21303910). Regulates transcription upon recruitment by YY1 to YY1-activated genes, where it acts as an essential coactivator (PubMed: 17721549). Involved in UV- damage excision DNA repair (PubMed: 20855601). The contribution to DNA double-strand break repair appears to be largely indirect through transcriptional regulation (PubMed: 20687897). Involved in DNA replication (PubMed: 20237820). Required for microtubule assembly during mitosis thereby regulating chromosome segregation cycle (PubMed: 20237820).
Cellular Location	Cytoplasm. Nucleus {ECO:0000255 PROSITE-ProRule:PRU00746, ECO:0000269 PubMed:16298340, ECO:0000269 PubMed:18026119, ECO:0000269 PubMed:20237820, ECO:0000269 PubMed:20971067}. Cytoplasm, cytoskeleton, spindle. Chromosome. Note=Localizes to the cytoplasm in quiescent cell (PubMed:20237820). Associates with spindle microtubules during mitosis (PubMed:20237820). Colocalizes with PCNA at replication forks during S-phase (PubMed:20237820). Recruited to DNA damage sites in a ACTR8-dependent manner (PubMed:20971067)
Tissue Location	According to PubMed:10574462, widely expressed. According to PubMed:16298340, specifically expressed in brain, liver and pancreas.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (INO80) Polyclonal Antibody, Unconjugated (AP56350) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.