10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TRIM45 Rabbit pAb

TRIM45 Rabbit pAb
Catalog # AP56381

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9H8W5
Reactivity	Rat, Human
Predicted	Pig, Dog, Mouse, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64359
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRIM45
Epitope Specificity	101-200/580
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the tripartite motif family. The encoded protein may function as a transcriptional repressor of the mitogen-activated protein kinase pathway. Alternatively spliced transcript variants have been described.[provided by RefSeq, Mar 2009]

Additional Information

Gene ID	80263
Other Names	E3 ubiquitin-protein ligase TRIM45, 2.3.2.27, RING finger protein 99, TRIM45, RNF99
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	TRIM45
Synonyms	RNF99

Function	E3 ubiquitin-protein ligase that plays a role in the regulation of inflammatory response (PubMed:36681779). Mechanistically, mediates the 'Lys-48'-linked polyubiquitination of TAB2, a regulatory protein of the kinase TAK1, leading to its degradation via the proteasomal pathway and inhibition of the TLR-mediated inflammatory immune response (PubMed:36681779). May act as a transcriptional repressor in mitogen-activated protein kinase signaling pathway.
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Expressed in skeletal muscle, brain, heart and pancreas.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: Siha Cell (Human) Lysate at 30 ug
Primary: Anti-TRIM45 (AP56381)at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 64kD
Observed band size: 64kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIM45) Polyclonal Antibody, Unconjugated (AP56381) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.