10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KLHDC1 Rabbit pAb

KLHDC1 Rabbit pAb
Catalog # AP56389

Product Information

Application	WB
Primary Accession	Q8N7A1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46720
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KLHDC1
Epitope Specificity	301-406/406
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	122773
Other Names	Kelch domain-containing protein 1, KLHDC1 {ECO:0000303|PubMed:16964437, ECO:0000312|HGNC:HGNC:19836}
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	KLHDC1 {ECO:0000303|PubMed:16964437, ECO:0000312|HGNC:HGNC:19836}
Function	Substrate-recognition component of a Cul5-RING (CRL5) E3 ubiquitin-protein ligase complex of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:32200094). The C-degron recognized by the DesCEND pathway is usually a motif of less than ten residues and can be present in full-length proteins, truncated

	proteins or proteolytically cleaved forms (PubMed:32200094). The CRL5(KLHDC1) complex mediates ubiquitination and degradation of truncated SELENOS selenoprotein produced by failed UGA/Sec decoding, which ends with a glycine (PubMed:32200094).
Cellular Location	Cytoplasm, cytosol
Tissue Location	Widely expressed, with high levels in skeletal muscle, pancreas and liver. Undetectable in peripheral blood leukocytes.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: HepG2 Cell (Human) Lysate at 40 ug
Primary: Anti-KLHDC1 (AP56389) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 47 kD
Observed band size: 47 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.