

KLHDC10 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56390

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession Popotivity	Dat Pavina
Heat	Ral, Buyine Dabbit
Clonality	Rabbit
Physical State	Liquid
	KI H conjugated synthetic pentide derived from human KI HDC10
Enitone Specificity	1_100/ <i>1</i> /2
Isotype	ΤσG
Purity	affinity purified by Protein Δ
T unity	annity partica by Hotem A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus
SIMILARITY	Contains 6 Kelch repeats.
SUBUNIT	Interacts with CUL2, TCEB1 and TCEB2; may be the substrate recognition
	component of an E3 ubiquitin ligase complex.
Important Note	Inis product as supplied is intended for research use only, not for use in
	numan, therapeutic or diagnostic applications.
Background Descriptions	KLHDC10 is a 442 amino acid protein that contains six keich repeats.
	two alternatively spliced isoforms. The gone encoding KLHDC10 maps to
	human chromosome 7 which houses over 1 000 genes and comprises nearly
	5% of the human genome. Chromosome 7 has been linked to Osteogenesis
	imperfecta. Pendred syndrome, Lissencephaly, Citrullinemia and
	Shwachman-Diamond syndrome. The deletion of a portion of the g arm of
	chromosome 7 is associated with Williams-Beuren syndrome, a condition
	characterized by mild mental retardation, an unusual comfort and
	friendliness with strangers and an elfin appearance. Deletions of portions of
	the q arm of chromosome 7 are also seen in a number of myeloid disorders,
	including cases of acute myelogenous leukemia and myelodysplasia.

Additional Information

Gene ID	23008
Other Names	Kelch domain-containing protein 10, KLHDC10, KIAA0265
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50 0,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KLHDC10 {ECO:0000303 PubMed:23102700, ECO:0000312 HGNC:HGNC:22194}
Function	Substrate-recognition component of a Cul2-RING (CRL2) E3 ubiquitin-protein ligase complex of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C- terminus of target proteins, leading to their ubiquitination and degradation (PubMed: <u>29779948</u> , PubMed: <u>33909987</u>). The C-degron recognized by the DesCEND pathway is usually a motif of less than ten residues and can be present in full-length proteins, truncated proteins or proteolytically cleaved forms (PubMed: <u>29779948</u> , PubMed: <u>33909987</u> , PubMed: <u>39548056</u>). The CRL2(KLHDC10) complex specifically recognizes proteins with a proline-glycine (Pro-Gly) or an alanine tail (CAT tail) at the C-terminus, leading to their ubiquitination and degradation (PubMed: <u>29779948</u> , PubMed: <u>33909987</u>). The CRL2(KLHDC10) complex is involved in the ribosome-associated quality control (RQC) pathway, which mediates the extraction of incompletely synthesized nascent chains from stalled ribosomes: CRL2(KLHDC10) acts downstream of NEMF and recognizes CAT tails associated with stalled nascent chains, leading to their ubiquitination and degradation (PubMed: <u>33909987</u>). Participates in the oxidative stress-induced cell death through MAP3K5 activation (PubMed: <u>23102700</u>). Inhibits PPP5C phosphatase activity on MAP3K5 (PubMed: <u>23102700</u>). Acts as a regulator of necroptosis (By similarity).
Cellular Location	Nucleus, Cytoplasm

Images



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KLHDC10) Polyclonal Antibody, Unconjugated (AP56390) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Sample: Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-KLHDC10(AP56390) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kD

Observed band size: 52 kD



Citations

• NEMF-mediated Listerin-independent mitochondrial translational surveillance by E3 ligase Pirh2 and mitochondrial protease ClpXP

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.