

KDEL Receptor Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56471

Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Bovine
Host
Clonality
Polyclonal
Calculated MW
P24390
Rat, Bovine
Rabbit
Polyclonal
24542

Additional Information

Gene ID 10945

Other Names ER lumen protein-retaining receptor 1, KDEL endoplasmic reticulum protein

retention receptor 1, KDEL receptor 1, Putative MAPK-activating protein PM23,

KDELR1, ERD2.1

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name KDELR1

Synonyms ERD2.1

Function Receptor for the C-terminal sequence motif K-D-E-L that is present on

endoplasmic reticulum resident proteins and that mediates their recycling

from the Golgi back to the endoplasmic reticulum.

Cellular Location Golgi apparatus membrane; Multi-pass membrane protein

{ECO:0000250 | UniProtKB:P33946}. Cytoplasmic vesicle, COPI-coated vesicle membrane; Multi-pass membrane protein {ECO:0000250 | UniProtKB:P33946}.

Endoplasmic reticulum membrane; Multi-pass membrane protein

{ECO:0000250 | UniProtKB:P33946}. Endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000250 | UniProtKB:P33946}; Multi-pass membrane protein {ECO:0000250 | UniProtKB:P33946} Note=Localized in the Golgi in the absence of bound proteins with the sequence motif K-D-E-L.

Trafficks back to the endoplasmic reticulum together with cargo proteins containing the sequence motif K-D-E-L $\,$

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.