

KIAA0907 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56497

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q7Z7F0
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64845
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KIAA0907
Epitope Specificity	451-550/614
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the UPF0469 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The KIAA0907 gene product has been provisionally designated KIAA0907 pending further characterization.

Additional Information

Gene ID	22889
Other Names	KH homology domain-containing protein 4 {ECO:0000312 HGNC:HGNC:29145}, Brings lots of money 7, Pre-mRNA splicing factor protein KHDC4, KHDC4 (HGNC:29145)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KHDC4 (HGNC:29145)
Function	RNA-binding protein involved in pre-mRNA splicing (PubMed: 19641227). Interacts with the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome (PubMed: 19641227). Involved in regulating splice site selection (PubMed: 19641227). Binds preferentially RNA with A/C rich sequences and poly-C stretches (PubMed: 23144703).
Cellular Location	Nucleus. Cytoplasm
Tissue Location	Ubiquitous. Expressed at high level in skeletal muscle, kidney, heart, brain and liver.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.