

KIAA1530 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56513

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q2YD98
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80591

Additional Information

Gene ID	57654
Other Names	UV-stimulated scaffold protein A, UVSSA, KIAA1530
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	UVSSA {ECO:0000303 PubMed:22466610, ECO:0000312 HGNC:HGNC:29304}
Function	Factor involved in transcription-coupled nucleotide excision repair (TC-NER), a mechanism that rapidly removes RNA polymerase II- blocking lesions from the transcribed strand of active genes (PubMed: 22466610 , PubMed: 22466611 , PubMed: 22466612 , PubMed: 32142649 , PubMed: 32355176 , PubMed: 34526721 , PubMed: 38316879 , PubMed: 38600235 , PubMed: 38600236). Acts as a key adapter that promotes recruitment of factors involved in TC-NER (PubMed: 22466611 , PubMed: 22466612 , PubMed: 32142649 , PubMed: 32355176 , PubMed: 34526721 , PubMed: 38600235 , PubMed: 38600236). Facilitates the ubiquitination of the elongating form of RNA polymerase II (RNA pol IIo) at DNA damage sites, thereby promoting RNA pol IIo backtracking and access by the TC-NER machinery to lesion sites (PubMed: 22466611 , PubMed: 32142649). Also promotes stabilization of ERCC6/CSB by recruiting deubiquitinating enzyme USP7 to TC-NER complexes, preventing UV-induced degradation of ERCC6 by the proteasome (PubMed: 22466611 , PubMed: 22466612). Mediates the recruitment of the TFIIH complex and other factors that are required for nucleotide excision repair to RNA polymerase II (PubMed: 32142649 ,

PubMed:[32355176](#), PubMed:[34526721](#), PubMed:[38600235](#), PubMed:[38600236](#)). Also required to inactivate stalled RNA polymerase II by blocking the access of TCEA1/TFIIS, thereby preventing reactivation of RNA polymerase II (PubMed:[38316879](#)). Not involved in processing oxidative damage (PubMed:[22466612](#)).

Cellular Location

Chromosome. Note=Accumulates at UV DNA damage sites

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.