

KIAA1731 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56519

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9C0D2
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	295176

Additional Information

Gene ID	85459
Other Names	Centrosomal protein of 295 kDa, CEP295 (HGNC:29366)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CEP295 (HGNC:29366)
Function	Centriole-enriched microtubule-binding protein involved in centriole biogenesis (PubMed: 20844083 , PubMed: 25131205 , PubMed: 27185865 , PubMed: 38154379). Essential for the generation of the distal portion of new-born centrioles in a CPAP- and CEP120-mediated elongation dependent manner during the cell cycle S/G2 phase after formation of the initiating cartwheel structure (PubMed: 27185865). Required for the recruitment of centriolar proteins, such as POC1B, POC5 and CEP135, into the distal portion of centrioles (PubMed: 27185865). Also required for centriole-to-centrosome conversion during mitotic progression, but is dispensable for cartwheel removal or centriole disengagement (PubMed: 25131205). Binds to and stabilizes centriolar microtubule (PubMed: 27185865). May be involved in ciliogenesis (PubMed: 38154379).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton. Note=Associates with both of the converted centrioles during G1 but becomes more enriched at the newly formed daughter (or unconverted) centrioles

during S, G2, and early M phases (PubMed:25131205, PubMed:32060285). In early S phase, localized at the procentriolar microtubule wall and enriched at the proximal ends of the centrioles in CPAP- and CEP135-dependent manner (PubMed:27185865). Colocalizes with SASS6 and CEP250 proteins (PubMed:25131205). Colocalizes with CEP135 and CEP192 at the centrosomes (PubMed:27185865). Associates with interphase microtubules and mitotic spindles (PubMed:27185865). Colocalizes with centriolar acetylated tubulin (PubMed:25131205).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.