

UCN2/Urocortin-2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56563

Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Dog
Host
Clonality
Polyclonal
Calculated MW
Physical State

Q96RP3
Rat, Dog
Rabbit
Polyclonal

Immunogen KLH conjugated synthetic peptide derived from human UCN2/Urocortin-2

Epitope Specificity 72-109/112

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Secreted

SIMILARITY Belongs to the sauvagine/corticotropin-releasing factor/urotensin I family.

SUBUNIT Binds with high affinity to CRF receptors 2-alpha and 2-beta.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This gene is a member of the sauvagine/corticotropin-releasing

factor/urotensin I family. It is structurally related to the

corticotropin-releasing factor (CRF) gene and the encoded product is an endogenous ligand for CRF type 2 receptors. In the brain it may be responsible for the effects of stress on appetite. In spite of the gene family

name similarity, the product of this gene has no sequence similarity to

urotensin II. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID 90226

Other Names Urocortin-2, Stresscopin-related peptide, Urocortin II, Ucn II,

Urocortin-related peptide, UCN2, SRP, URP

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

UCN2 Name

SRP, URP **Synonyms**

Function

Suppresses food intake, delays gastric emptying and decreases heat-induced edema. Might represent an endogenous ligand for maintaining homeostasis

after stress.

Secreted. **Cellular Location**

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.