

TRIT1 Rabbit pAb

TRIT1 Rabbit pAb
Catalog # AP56571

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9H3H1
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Rabbit, Chicken, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52725
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRIT1
Epitope Specificity	121-220/467
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm and Mitochondrion.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Catalyzes the transfer of a dimethylallyl group onto the adenine at position 37 of both cytosolic and mitochondrial tRNAs, leading to the formation of N6-(dimethylallyl)adenosine (i(6)A).

Additional Information

Gene ID	54802
Other Names	tRNA dimethylallyltransferase, 2.5.1.75, Isopentenyl-diphosphate:tRNA isopentenyltransferase, IPP transferase, IPPT, hGRO1, tRNA isopentenyltransferase 1, IPTase, TRIT1, IPT, MOD5
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TRIT1
Synonyms	IPT, MOD5


Function Catalyzes the transfer of a dimethylallyl group onto the adenine at position 37 of both cytosolic and mitochondrial tRNAs, leading to the formation of N6-(dimethylallyl)adenosine (i6A37) (PubMed:[11111046](#), PubMed:[24126054](#), PubMed:[24901367](#), PubMed:[34774131](#)). Mediates modification of a limited subset of tRNAs: tRNA(Ser)(AGA), tRNA(Ser)(CGA), tRNA(Ser)(UGA), as well as partial modification of the selenocysteine tRNA(Ser)(UCA) (PubMed:[24126054](#)). TRIT1 is therefore required for selenoprotein expression (PubMed:[24126054](#)).

Cellular Location [Isoform 1]: Mitochondrion [Isoform 2]: Cytoplasm. [Isoform 5]: Cytoplasm.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: Uterus(Mouse) Lysate at 40 ug
Primary: Anti-TRIT1(AP56571)at 1/300 dilution
Secondary: IRDye800CW Goat Anti-RabbitIgG at 1/20000 dilution
Predicted band size: 48kD
Observed band size: 53kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIT1) Polyclonal Antibody, Unconjugated (AP56571) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.