

# TPMT Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56588

## Product Information

<b>Application</b>	IHC-P, IHC-F, IF, ICC, E
<b>Primary Accession</b>	<a href="#">P51580</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	28180

## Additional Information

<b>Gene ID</b>	7172
<b>Other Names</b>	Thiopurine S-methyltransferase, 2.1.1.67, Thiopurine methyltransferase, TPMT
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

<b>Name</b>	TPMT
<b>Function</b>	Catalyzes the S-methylation of thiopurine drugs such as 6- mercaptopurine (also called mercaptopurine, 6-MP or its brand name Purinethol) and 6-thioguanine (also called tioguanine or 6-TG) using S- adenosyl-L-methionine as the methyl donor (PubMed: <a href="#">18484748</a> , PubMed: <a href="#">657528</a> ). TPMT activity modulates the cytotoxic effects of thiopurine prodrugs. A natural substrate for this enzyme has yet to be identified.
<b>Cellular Location</b>	Cytoplasm.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.