

TPRKB Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56590

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9Y3C4
Reactivity	Rat
Predicted	Human, Mouse, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19661
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TPRKB
Epitope Specificity	1-100/175
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
SIMILARITY	Belongs to the CGI121/TPRKB family.
SUBUNIT	Component of the EKC/KEOPS complex composed of at least TP53RK, TPRKB, OSGEP and LAGE3; the whole complex dimerizes. Interacts with TP53RK/PRPK.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


Additional Information

Gene ID	51002
Other Names	EKC/KEOPS complex subunit TPRKB, PRPK-binding protein, TP53RK-binding protein, TPRKB (HGNC:24259)
Target/Specificity	Widely expressed.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TPRKB (HGNC:24259)
Function	Component of the EKC/KEOPS complex that is required for the formation of a threonylcarbamoyl group on adenosine at position 37 (t(6)A37) in tRNAs that read codons beginning with adenine (PubMed: 22912744 , PubMed: 28805828). The complex is probably involved in the transfer of the threonylcarbamoyl moiety of threonylcarbamoyl-AMP (TC-AMP) to the N6 group of A37 (PubMed: 22912744 , PubMed: 28805828). TPRKB acts as an allosteric effector that regulates the t(6)A activity of the complex. TPRKB is not required for tRNA modification (PubMed: 22912744 , PubMed: 28805828).
Cellular Location	Cytoplasm, cytosol. Nucleus
Tissue Location	Widely expressed..

Images



Paraformaldehyde-fixed, paraffin embedded (rat uterus tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TPRKB) Polyclonal Antibody, Unconjugated (AP56590) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.