

JAML Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56603

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession

Reactivity

Host

Clonality

Calculated MW

Q86YT9

Human

Rabbit

Polyclonal

44339

Additional Information

Gene ID 120425

Other Names Junctional adhesion molecule-like {ECO:0000312 | HGNC:HGNC:19084},

Adhesion molecule interacting with CXADR antigen 1, Dendritic cell-specific

protein CREA7-1, JAML (HGNC:19084), AMICA1

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name JAML (HGNC:19084)

Synonyms AMICA1

Function Transmembrane protein of the plasma membrane of leukocytes that control

their migration and activation through interaction with CXADR, a plasma membrane receptor found on adjacent epithelial and endothelial cells. The interaction between both receptors mediates the activation of gamma-delta T-cells, a subpopulation of T-cells residing in epithelia and involved in tissue homeostasis and repair. Upon epithelial CXADR-binding, JAML induces downstream cell signaling events in gamma-delta T-cells through PI3-kinase and MAP kinases. It results in proliferation and production of cytokines and growth factors by T- cells that in turn stimulate epithelial tissues repair. It also controls the transmigration of leukocytes within epithelial and endothelial tissues through adhesive interactions with epithelial and endothelial CXADR.

Cellular Location Cell membrane; Single-pass type I membrane protein. Cell junction.

Note=Localized at the plasma membrane and enriched in areas of cell-cell

contacts (PubMed:12869515)

Tissue Location

Expression is restricted to the hematopoietic tissues with the exception of liver. Expressed in fetal liver, spleen and thymus. Preferentially expressed by mature leukocytes (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.