

KAAG1 Rabbit pAb

KAAG1 Rabbit pAb
Catalog # AP56611

Product Information

Primary Accession	Q9UBP8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	8969
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KAAG1
Epitope Specificity	31-84/84
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information


Other Names	Kidney-associated antigen 1, RU2 antisense gene protein, KAAG1, RU2AS
Dilution	Flow-Cyt=1ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KAAG1
Synonyms	RU2AS
Tissue Location	Expressed in testis and kidney, and, at lower levels, in urinary bladder and liver. Expressed by a high proportion of tumors of various histologic origin, including melanomas, sarcomas and colorectal carcinomas.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Blank control: A549.

Primary Antibody (green line): Rabbit Anti-KAAG1 antibody (AP56611)

Dilution: 1 μg / 10^6 cells;


Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 μg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: A549.

Primary Antibody (green line): Rabbit Anti-KAAG1 antibody (AP56611)

Dilution: 1 μg / 10^6 cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 μg /test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.