

TOM1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56615

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O60784
Reactivity	Rat, Pig, Dog, Chimpanzee, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53818

Additional Information

Gene ID	10043
Other Names	Target of Myb protein 1, TOM1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TOM1 (HGNC:11982)
Function	Adapter protein that plays a role in the intracellular membrane trafficking of ubiquitinated proteins, thereby participating in autophagy, ubiquitination-dependent signaling and receptor recycling pathways (PubMed: 14563850 , PubMed: 15047686 , PubMed: 23023224 , PubMed: 25588840 , PubMed: 26320582 , PubMed: 31371777). Acts as a MYO6/Myosin VI adapter protein that targets MYO6 to endocytic structures (PubMed: 23023224). Together with MYO6, required for autophagosomal delivery of endocytic cargo, the maturation of autophagosomes and their fusion with lysosomes (PubMed: 23023224). MYO6 links TOM1 with autophagy receptors, such as TAX1BP1; CALCOCO2/NDP52 and OPTN (PubMed: 31371777). Binds to polyubiquitinated proteins via its GAT domain (PubMed: 14563850). In a complex with TOLLIP, recruits ubiquitin-conjugated proteins onto early endosomes (PubMed: 15047686). The Tom1-Tollip complex may regulate endosomal trafficking by linking polyubiquitinated proteins to clathrin (PubMed: 14563850 , PubMed: 15047686). Mediates clathrin recruitment to early endosomes by ZFYVE16 (PubMed: 15657082). Modulates

binding of TOLLIP to phosphatidylinositol 3-phosphate (PtdIns(3)P) via binding competition; the association with TOLLIP may favor the release of TOLLIP from endosomal membranes, allowing TOLLIP to commit to cargo trafficking (PubMed:[26320582](#)). Acts as a phosphatidylinositol 5-phosphate (PtdIns(5)P) effector by binding to PtdIns(5)P, thereby regulating endosomal maturation (PubMed:[25588840](#)). PtdIns(5)P-dependent recruitment to signaling endosomes may block endosomal maturation (PubMed:[25588840](#)). Also inhibits Toll-like receptor (TLR) signaling and participates in immune receptor recycling (PubMed:[15047686](#), PubMed:[26320582](#)).

Cellular Location

Cytoplasm. Endosome membrane; Peripheral membrane protein. Early endosome membrane; Peripheral membrane protein. Note=Localized to endo/exosomal vesicles (PubMed:31263572). Enriched on signaling endosomes (PubMed:25588840). Recruited to early endosomes by TOLLIP and by PtdIns(5)P (PubMed:15047686, PubMed:25588840, PubMed:26320582)

Tissue Location

Widely expressed. Highly expressed in skeletal muscle, heart, placenta and liver.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.