

SCD5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56640

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q86SK9
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37610
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SCD5
Epitope Specificity	251-330/330
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum membrane.
SIMILARITY	Belongs to the fatty acid desaturase family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Stearoyl-CoA desaturase (SCD; EC 1.14.99.5) is an integral membrane protein of the endoplasmic reticulum that catalyzes the formation of monounsaturated fatty acids from saturated fatty acids. SCD may be a key regulator of energy metabolism with a role in obesity and dislipidemia. Four SCD isoforms, Scd1 through Scd4, have been identified in mouse. In contrast, only 2 SCD isoforms, SCD1 (MIM 604031) and SCD5, have been identified in human. SCD1 shares about 85% amino acid identity with all 4 mouse SCD isoforms, as well as with rat Scd1 and Scd2. In contrast, SCD5 shares limited homology with the rodent SCDs and appears to be unique to primates (Wang et al., 2005 [PubMed 15907797]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	79966
Other Names	Stearoyl-CoA desaturase 5, 1.14.19.1, Acyl-CoA-desaturase 4, HSCD5, Stearoyl-CoA 9-desaturase, Stearoyl-CoA desaturase 2, SCD5
Target/Specificity	Detected in fetal brain, and at lower levels in fetal kidney. Detected in adult brain and pancreas, and at lower levels in kidney and lung.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	SCD5
Function	Stearoyl-CoA desaturase that utilizes O(2) and electrons from reduced cytochrome b5 to introduce the first double bond into saturated fatty acyl-CoA substrates. Catalyzes the insertion of a cis double bond at the delta-9 position into fatty acyl-CoA substrates including palmitoyl-CoA and stearoyl-CoA (PubMed: 15610069 , PubMed: 15907797 , PubMed: 22745828). Gives rise to a mixture of 16:1 and 18:1 unsaturated fatty acids (PubMed: 15610069 , PubMed: 15907797). Involved in neuronal cell proliferation and differentiation through down-regulation of EGFR/AKT/MAPK and Wnt signaling pathways (PubMed: 22745828).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein
Tissue Location	Detected in fetal brain, and at lower levels in fetal kidney. Detected in adult brain and pancreas, and at lower levels in kidney and lung. Expressed in spiral ganglion cells and the organ of Corti of fetal cochlea (PubMed:31972369).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.