

hnRNP H Rabbit pAb

hnRNP H Rabbit pAb Catalog # AP56661

Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession <u>P31943</u>

Predicted Human, Mouse, Rat, Zebrafish, Rhesus, Gorilla, Opossum, Orangutan,

Elephant

Host Rabbit
Clonality Polyclonal
Calculated MW 49229
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human hnRNP H

Epitope Specificity 151-250/449

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus; nucleoplasm.

SIMILARITY Contains 3 RRM (RNA recognition motif) domains.

SUBUNITPart of a ternary complex containing FUBP2, PTBP1, PTBP2 and HNRNPH1.
Identified in the spliceosome C complex. Interacts with IGF2BP1, Interacts

Identified in the spliceosome C complex. Interacts with IGF2BP1. Interacts with CUGBP1; the interaction is RNA-dependent. Interacts with MBNL1; the

interaction in RNA-independent.

Important NoteThis product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This gene encodes a member of a subfamily of ubiquitously expressed

heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins that complex with heterogeneous nuclear RNA. These proteins are associated with pre-mRNAs in the nucleus and appear to

influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some may shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has three repeats of quasi-RRM domains that bind to RNA and is very similar to the family member HNRPF. This gene may be associated with hereditary lymphedema type I. Alternatively spliced transcript variants have been

described [provided by RefSeq, Mar 2012]

Additional Information

Gene ID 3187

Other Names Heterogeneous nuclear ribonucleoprotein H, hnRNP H, Heterogeneous

nuclear ribonucleoprotein H, N-terminally processed, HNRNPH1, HNRPH,

HNRPH1

Target/Specificity Expressed ubiquitously.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-

500,ELISA=1:5000-10000

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name HNRNPH1

Synonyms HNRPH, HNRPH1

Function This protein is a component of the heterogeneous nuclear ribonucleoprotein

(hnRNP) complexes which provide the substrate for the processing events that pre-mRNAs undergo before becoming functional, translatable mRNAs in the cytoplasm. Mediates pre-mRNA alternative splicing regulation. Inhibits, together with CUGBP1, insulin receptor (IR) pre-mRNA exon 11 inclusion in

myoblast. Binds to the IR RNA. Binds poly(RG).

Cellular Location Nucleus, nucleoplasm.

Tissue Location Expressed ubiquitously.

Background

This gene encodes a member of a subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins that complex with heterogeneous nuclear RNA. These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some may shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has three repeats of quasi-RRM domains that bind to RNA and is very similar to the family member HNRPF. This gene may be associated with hereditary lymphedema type I. Alternatively spliced transcript variants have been described [provided by RefSeq, Mar 2012]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.