

## **HOXB7** Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56673

## **Product Information**

**Application** WB, IHC-P, IHC-F, IF, ICC, E

**Primary Accession** P09629 Reactivity Rat Host Rabbit Clonality Polyclonal **Calculated MW** 24015 **Physical State** Liquid

**Immunogen** KLH conjugated synthetic peptide derived from human HOXB7

101-200/217 **Epitope Specificity** 

Isotype IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION

**SIMILARITY** 

Belongs to the Antp homeobox family. Contains 1 homeobox DNA-binding

domain.

This product as supplied is intended for research use only, not for use in **Important Note** 

human, therapeutic or diagnostic applications.

**Background Descriptions** The Hox proteins play a role in development and cellular differentiation by

regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. The mammalian HOX gene complex consists of 39 genes that are located on four linkage groups, which are dispersed over four chromosomes. A segment of the HoxB7 proximal promoter drives renal expression of reporter genes specifically in the ureteric bud and collecting ducts. Expression levels of HoxB7 are lower in lymph node metastasis-positive cancer tissues than negative cancer tissues. These results suggest that aberrant expression of HOX genes is related to the development of breast cancer and malignant

behavior of cancer cells.

## **Additional Information**

Gene ID 3217

**Other Names** Homeobox protein Hox-B7, Homeobox protein HHO.C1, Homeobox protein

Hox-2C, HOXB7, HOX2C

WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50 Dilution

0,ELISA=1:5000-10000

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce **Format** 

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Protein Information**

Name HOXB7

Synonyms HOX2C

**Function** Sequence-specific transcription factor which is part of a developmental

regulatory system that provides cells with specific positional identities on the

anterior-posterior axis.

Cellular Location Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.