

OSBPL5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56728

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q9H0X9
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98616
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human OSBPL5
Epitope Specificity	51-150/879
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors that play a key role in the maintenance of cholesterol balance in the body. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain. This gene has been shown to be imprinted, with preferential expression from the maternal allele only in placenta. Transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2010]


Additional Information

Gene ID	114879
Other Names	Oxysterol-binding protein-related protein 5, ORP-5, OSBP-related protein 5, Oxysterol-binding protein homolog 1, OSBPL5, KIAA1534, OBPH1 {ECO:0000303 PubMed:12504849}, ORP5
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	OSBPL5
Synonyms	KIAA1534, OBPH1 {ECO:0000303 PubMed:1250
Function	Lipid transporter involved in lipid countertransport between the endoplasmic reticulum and the plasma membrane: specifically exchanges phosphatidylserine with phosphatidylinositol 4-phosphate (PI4P), delivering phosphatidylserine to the plasma membrane in exchange for PI4P, which is degraded by the SAC1/SACM1L phosphatase in the endoplasmic reticulum. Binds phosphatidylserine and PI4P in a mutually exclusive manner (PubMed: 23934110 , PubMed: 26206935). May cooperate with NPC1 to mediate the exit of cholesterol from endosomes/lysosomes (PubMed: 21220512). Binds 25-hydroxycholesterol and cholesterol (PubMed: 17428193).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein. Note=Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Localizes to the cortical endoplasmic reticulum at the EPCS
Tissue Location	Ubiquitously expressed.

Images



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OSBPL5) Polyclonal Antibody, Unconjugated (AP56728) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OSBPL5) Polyclonal Antibody, Unconjugated (AP56728) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.