

OSBPL5 Rabbit pAb

OSBPL5 Rabbit pAb

Catalog # AP56728

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9H0X9
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98616
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human OSBPL5
Epitope Specificity	51-150/879
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the oxysterol-binding protein (OSBP) family, a group of intracellular lipid receptors that play a key role in the maintenance of cholesterol balance in the body. Most members contain an N-terminal pleckstrin homology domain and a highly conserved C-terminal OSBP-like sterol-binding domain. This gene has been shown to be imprinted, with preferential expression from the maternal allele only in placenta. Transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2010]

Additional Information

Gene ID	114879
Other Names	Oxysterol-binding protein-related protein 5, ORP-5, OSBP-related protein 5, Oxysterol-binding protein homolog 1, OSBPL5, KIAA1534, OBPH1 {ECO:0000303 PubMed:12504849}, ORP5
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	OSBPL5 (HGNC:16392)
Function	Lipid transporter involved in lipid countertransport between the endoplasmic reticulum and the plasma membrane: specifically exchanges phosphatidylserine with phosphatidylinositol 4-phosphate (PI4P), delivering phosphatidylserine to the plasma membrane in exchange for PI4P, which is degraded by the SAC1/SACM1L phosphatase in the endoplasmic reticulum. Binds phosphatidylserine and PI4P in a mutually exclusive manner (PubMed: 23934110 , PubMed: 26206935). May cooperate with NPC1 to mediate the exit of cholesterol from endosomes/lysosomes (PubMed: 21220512). Binds 25-hydroxycholesterol and cholesterol (PubMed: 17428193).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein. Note=Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Localizes to the cortical endoplasmic reticulum at the EPCS
Tissue Location	Ubiquitously expressed.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OSBPL5) Polyclonal Antibody, Unconjugated (AP56728) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.