

FGFBP1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56797

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q14512
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26264
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FGFBP
Epitope Specificity	151-234/234
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted, extracellular space. Cell membrane; Peripheral membrane protein. Note=Extracellular and plasma membrane-associated. Colocalizes with HSPG2 in the pericellular environment of squamous cell carcinomas.
SIMILARITY	Belongs to the fibroblast growth factor-binding protein family.
SUBUNIT	Found in a complex with FGFBP1, FGF1 and FGF2. Interacts with FGF1, FGF2, FGF7, FGF10, FGF22 and HSPG2.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a secreted fibroblast growth factor carrier protein. The encoded protein plays a critical role in cell proliferation, differentiation and migration by binding to fibroblast growth factors and potentiating their biological effects on target cells. The encoded protein may also play a role in tumor growth as an angiogenic switch molecule, and expression of this gene has been associated with several types of cancer including pancreatic and colorectal adenocarcinoma. A pseudogene of this gene is also located on the short arm of chromosome 4. [provided by RefSeq, Nov 2011]

Additional Information

Gene ID	9982
Other Names	Fibroblast growth factor-binding protein 1, FGF-BP, FGF-BP1, FGF-binding protein 1, FGFBP-1, 17 kDa heparin-binding growth factor-binding protein, 17 kDa HBGF-binding protein, HBp17, FGFBP1, FGFBP, HBP17
Target/Specificity	Expressed in the suprabasal region of the epidermis, in hair follicles, the basement membrane at the dermo-epidermal junction (occasionally extending into the basement membrane of dermal blood vessels), wounded skin and several invasive squamous cell carcinomas (at protein level).


Expressed in normal and wounded skin and various squamous cell carcinomas.

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	FGFBP1
Synonyms	FGFBP, HBP17
Function	Acts as a carrier protein that release fibroblast-binding factors (FGFs) from the extracellular matrix (EM) storage and thus enhance the mitogenic activity of FGFs. Enhances FGF2 signaling during tissue repair, angiogenesis and in tumor growth.
Cellular Location	Secreted, extracellular space. Cell membrane; Peripheral membrane protein. Note=Extracellular and plasma membrane- associated. Colocalizes with HSPG2 in the pericellular environment of squamous cell carcinomas.
Tissue Location	Expressed in the suprabasal region of the epidermis, in hair follicles, the basement membrane at the dermo- epidermal junction (occasionally extending into the basement membrane of dermal blood vessels), wounded skin and several invasive squamous cell carcinomas (at protein level). Expressed in normal and wounded skin and various squamous cell carcinomas

Images




Sample:
A431(Human) Cell Lysate at 30 ug
Primary: Anti-FGFBP1 (AP56797) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 28 kD
Observed band size: 50 kD


Paraformaldehyde-fixed, paraffin embedded (human skin cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FGFBP1) Polyclonal Antibody, Unconjugated (AP56797) at 1:400 overnight at 4°C,



followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: mouse intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-FGFBP1 Polyclonal Antibody, Unconjugated(AP56797) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse stomach tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-FGFBP1 Polyclonal Antibody, Unconjugated(AP56797) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.