

HSPB4/Alpha A Crystallin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56803

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	P02489
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19909

Additional Information

Gene ID	102724652;1409
Other Names	Alpha-crystallin A chain, Heat shock protein beta-4, HspB4, Alpha-crystallin A(1-172), Alpha-crystallin A(1-168), Alpha-crystallin A(1-162), CRYAA, CRYA1, HSPB4
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CRYAA
Synonyms	CRYA1, HSPB4
Function	Contributes to the transparency and refractive index of the lens (PubMed: 18302245). In its oxidized form (absence of intramolecular disulfide bond), acts as a chaperone, preventing aggregation of various proteins under a wide range of stress conditions (PubMed: 18199971 , PubMed: 19595763 , PubMed: 22120592 , PubMed: 31792453). Required for the correct formation of lens intermediate filaments as part of a complex composed of BFSP1, BFSP2 and CRYAA (PubMed: 28935373).
Cellular Location	Cytoplasm. Nucleus. Note=Translocates to the nucleus during heat shock and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles

Tissue Location

Expressed in the eye lens (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.