

## Taxilin alpha Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56807

## **Product Information**

**Application** IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Rat
Host
Clonality
Polyclonal
Calculated MW
Physical State
P40222
Rat
Rabbit
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived from human IL-14

Epitope Specificity 401-500/546

Isotype IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SIMILARITY** Belongs to the taxilin family.

**SUBUNIT** Binds to the C-terminal coiled coil region of syntaxin family members STX1A,

STX3A and STX4A, but not when these proteins are complexed with SNAP25, VAMP2 or STXBP1, suggesting that it interacts with syntaxins that do not form

the SNARE complex.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** Alpha Taxilin may be involved in intracellular vesicle traffic and potentially in

calcium dependent exocytosis in neuroendocrine cells. It binds to the C terminal coiled coil region of syntaxin family members STX1A, STX3A and STX4A, but not when these proteins are complexed with SNAP25, VAMP2 or STXBP1, suggesting that it interacts with syntaxins that do not form the SNARE

complex.

## **Additional Information**

**Gene ID** 200081

Other Names Alpha-taxilin, TXLNA, TXLN

**Target/Specificity** Ubiquitous, with much higher expression in heart, kidney, liver and pancreas.

**Dilution** IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Protein Information**

**TXLNA** Name

**Synonyms TXLN** 

May be involved in intracellular vesicle traffic and potentially in calcium-dependent exocytosis in neuroendocrine cells. **Function** 

Ubiquitous, with much higher expression in heart, kidney, liver and pancreas. **Tissue Location** 

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.