

MOSC2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56809

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q969Z3
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38023

Additional Information

Gene ID	54996
Other Names	Mitochondrial amidoxime reducing component 2, mARC2, 1.7.-., Molybdenum cofactor sulfurase C-terminal domain-containing protein 2, MOSC domain-containing protein 2, Moco sulfurase C-terminal domain-containing protein 2, MTARC2 (HGNC:26064), MARC2, MOSC2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MTARC2 (HGNC:26064)
Synonyms	MARC2, MOSC2
Function	Catalyzes the reduction of N-oxygenated molecules, acting as a counterpart of cytochrome P450 and flavin-containing monooxygenases in metabolic cycles (PubMed: 21029045 , PubMed: 24423752). As a component of prodrug-converting system, reduces a multitude of N-hydroxylated prodrugs particularly amidoximes, leading to increased drug bioavailability (PubMed: 21029045 , PubMed: 24423752). May be involved in mitochondrial N(omega)-hydroxy-L-arginine (NOHA) reduction, regulating endogenous nitric oxide levels and biosynthesis (PubMed: 21029045). Postulated to cleave the N-OH bond of N-hydroxylated substrates in concert with electron transfer from NADH to cytochrome b5 reductase then to cytochrome b5, the ultimate electron donor that primes the active site for substrate reduction (PubMed: 21029045).

Cellular Location

Mitochondrion outer membrane; Peripheral membrane protein. Peroxisome

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.