

MSH4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56866

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O15457
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	104756
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MSH4
Epitope Specificity	771-870/936
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear
SIMILARITY	Belongs to the DNA mismatch repair MutS family.
SUBUNIT	Heterooligomer of MSH4 and MSH5.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the DNA mismatch repair mutS family. This member is a meiosis-specific protein that is not involved in DNA mismatch correction, but is required for reciprocal recombination and proper segregation of homologous chromosomes at meiosis I. This protein and MSH5 form a heterodimer which binds uniquely to a Holliday Junction and its developmental progenitor, thus provoking ADP-ATP exchange, and stabilizing the interaction between parental chromosomes during meiosis double-stranded break repair. [provided by RefSeq, Aug 2011]

Additional Information


Gene ID	4438
Other Names	MutS protein homolog 4, hMSH4, MSH4
Target/Specificity	Testis and ovary specific.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	MSH4
Function	Involved in meiotic recombination. Required for reciprocal recombination and proper segregation of homologous chromosomes at meiosis.
Cellular Location	Chromosome {ECO:0000250 UniProtKB:Q99MT2}.
Tissue Location	Highly expressed in testis (PubMed:33437391, PubMed:9299235). Also expressed in the ovary (PubMed:9299235)

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MSH4) Polyclonal Antibody, Unconjugated (AP56866) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MSH4) Polyclonal Antibody, Unconjugated (AP56866) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.