

MSL3L1 Rabbit pAb

MSL3L1 Rabbit pAb

Catalog # AP56870

Product Information

Application	WB
Primary Accession	Q8N5Y2
Reactivity	Human
Predicted	Pig, Dog, Mouse, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59824
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MSL3L1
Epitope Specificity	1-100/521
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a nuclear protein that is similar to the product of the Drosophila male-specific lethal-3 gene. The Drosophila protein plays a critical role in a dosage-compensation pathway, which equalizes X-linked gene expression in males and females. Thus, the human protein is thought to play a similar function in chromatin remodeling and transcriptional regulation, and it has been found as part of a complex that is responsible for histone H4 lysine-16 acetylation. This gene can undergo X inactivation. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 2, 7 and 8. [provided by RefSeq, Jul 2010]

Additional Information

Gene ID	10943
Other Names	MSL complex subunit 3 {ECO:0000312 HGNC:HGNC:7370}, Male-specific lethal 3 homolog, Male-specific lethal-3 homolog 1, Male-specific lethal-3 protein-like 1, MSL3-like 1, MSL3 {ECO:0000303 PubMed:16227571, ECO:0000312 HGNC:HGNC:7370}
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	MSL3 {ECO:0000303 PubMed:16227571, ECO:0000312 HGNC:HGNC:7370}
Function	Non-catalytic component of the MSL histone acetyltransferase complex, a multiprotein complex that mediates the majority of histone H4 acetylation at 'Lys-16' (H4K16ac), an epigenetic mark that prevents chromatin compaction (PubMed:16227571, PubMed:16543150, PubMed:20018852, PubMed:20657587, PubMed:20943666, PubMed:21217699, PubMed:30224647, PubMed:33837287). The MSL complex is required for chromosome stability and genome integrity by maintaining homeostatic levels of H4K16ac (PubMed:33837287). The MSL complex is also involved in gene dosage by promoting up-regulation of genes expressed by the X chromosome (By similarity). X up-regulation is required to compensate for autosomal biallelic expression (By similarity). The MSL complex also participates in gene dosage compensation by promoting expression of Tsix non-coding RNA (By similarity). Acts as a histone reader that specifically recognizes and binds histone H4 monomethylated at 'Lys-20' (H4K20Me1) in a DNA-dependent manner and is proposed to be involved in chromosomal targeting of the MSL complex (PubMed:20657587, PubMed:20943666). May play a role X inactivation in females (PubMed:21217699).
Cellular Location	Nucleus.
Tissue Location	Expressed in many tissues including liver, pancreas, heart, lung, kidney, skeletal muscle, brain, and placenta, with highest expression in skeletal muscle and heart

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.