

mu Crystallin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56891

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q14894
Reactivity	Rat, Pig, Cat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33776

Additional Information

Gene ID	1428
Other Names	Ketimine reductase mu-crystallin, 1.5.1.25, NADP-regulated thyroid-hormone-binding protein, CRYM, THBP
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CRYM (HGNC:2418)
Function	Catalyzes the NAD(P)H-dependent reduction of imine double bonds of a number of cyclic ketimine substrates, including sulfur- containing cyclic ketimines (PubMed: 21332720 , PubMed: 25931162). Under physiological conditions, it efficiently catalyzes delta(1)- piperideine-2-carboxylate (P2C) and delta(1)-pyrroline-2-carboxylate (Pyr2C) reduction, suggesting a central role in lysine and glutamate metabolism (PubMed: 25931162). Additional substrates are delta(2)- thiazoline-2-carboxylate (T2C), 3,4-dehydrothiomorpholine-3-carboxylate (AECK), and (R)-lanthionine ketimine (LK) that is reduced at very low rate compared to other substrates (PubMed: 25931162). Also catalyzes the NAD(P)H-dependent reduction of (S)-cystathionine ketimine (CysK) (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in neural tissues, muscle and kidney (PubMed:1384048). Expressed in the inner ear (PubMed:12471561)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.