

HspBP1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56944

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NZL4
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39303
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HspBP1
Epitope Specificity	51-150/362
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 4 ARM repeats.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Hsp70-interacting protein (HspBP1) belongs to a family of eukaryotic proteins identified as nucleotide exchange factors for Hsp70, which exhibit varying degrees of compartment and species specificity. HspBP1 interferes with the CHIP-induced degradation of immature forms of the cystic fibrosis transmembrane conductance regulator (CFTR) and stimulates CFTR maturation. HspBP1 binds to Hsp70, inhibits its activity and promotes dissociation of nucleotides from the Hsp70 ATPase domain. It is a protein mainly expressed in heart and skeletal muscle.

Additional Information


Gene ID	23640
Other Names	Hsp70-binding protein 1, HspBP1, Heat shock protein-binding protein 1, Hsp70-binding protein 2, HspBP2, Hsp70-interacting protein 1, Hsp70-interacting protein 2, HSPBP1 (HGNC:24989), HSPBP
Target/Specificity	Ubiquitous.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HSPBP1 (HGNC:24989)
Synonyms	HSPBP
Function	Inhibits HSPA1A chaperone activity by changing the conformation of the ATP-binding domain of HSPA1A and interfering with ATP binding. Interferes with ubiquitination mediated by STUB1 and inhibits chaperone-assisted degradation of immature CFTR.
Tissue Location	Ubiquitous..

Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HspBP1) Polyclonal Antibody, Unconjugated (AP56944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.