

hUPF2/RENT2 Rabbit pAb

hUPF2/RENT2 Rabbit pAb

Catalog # AP56951

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9HAU5
Reactivity	Mouse
Predicted	Rat, Dog, Gorilla, Human, Rabbit, Sheep, Opossum
Host	Rabbit
Clonality	Polyclonal
Calculated MW	147810
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human hUPF2/RENT2
Epitope Specificity	951-1050/1272
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm > perinuclear region.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located in the perinuclear area. It interacts with translation release factors and the proteins that are functional homologs of yeast Upf1p and Upf3p. Two splice variants have been found for this gene; both variants encode the same protein. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	26019
Other Names	Regulator of nonsense transcripts 2 {ECO:0000312 HGNC:HGNC:17854}, Up-frameshift suppressor 2 homolog, hUpf2, UPF2 (HGNC:17854)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	UPF2 (HGNC:17854)
Function	Involved in nonsense-mediated decay (NMD) of mRNAs containing premature stop codons by associating with the nuclear exon junction complex (EJC). Recruited by UPF3B associated with the EJC core at the cytoplasmic side of the nuclear envelope and the subsequent formation of an UPF1-UPF2-UPF3 surveillance complex (including UPF1 bound to release factors at the stalled ribosome) is believed to activate NMD. In cooperation with UPF3B stimulates both ATPase and RNA helicase activities of UPF1. Binds spliced mRNA.
Cellular Location	Cytoplasm, perinuclear region. Cytoplasm {ECO:0000250 UniProtKB:A2AT37}
Tissue Location	Ubiquitous..

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (hUPF2) Polyclonal Antibody, Unconjugated (AP56951) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.