

TNPO3 Rabbit pAb

TNPO3 Rabbit pAb
Catalog # AP56966

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9Y5L0
Predicted	Rat, Human, Mouse, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	104203
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TNPO3
Epitope Specificity	821-923/923
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a nuclear import receptor for serine/arginine-rich (SR) proteins such as the splicing factors SFRS1 and SFRS2. The encoded protein has also been shown to be involved in HIV-1 infection, apparently through interaction with the HIV-1 capsid protein. Two transcript variants encoding different isoforms as well as a noncoding transcript have been found for this gene.[provided by RefSeq, Jul 2010]

Additional Information

Gene ID	23534
Other Names	Transportin-3, Importin-12, Imp12, Transportin-SR, TRN-SR, TNPO3 {ECO:0000303 PubMed:23667635, ECO:0000312 HGNC:HGNC:17103}
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TNPO3 {ECO:0000303 PubMed:23667635,
-------------	---------------------------------------

Function

Importin, which transports target proteins into the nucleus (PubMed:[10366588](#), PubMed:[10713112](#), PubMed:[11517331](#), PubMed:[12628928](#), PubMed:[24449914](#)). Specifically mediates the nuclear import of splicing factor serine/arginine (SR) proteins, such as RBM4, SFRS1 and SFRS2, by recognizing phosphorylated SR domains (PubMed:[10366588](#), PubMed:[10713112](#), PubMed:[11517331](#), PubMed:[12628928](#), PubMed:[24449914](#)). Also mediates the nuclear import of serine/arginine (SR) protein CPSF6, independently of CPSF6 phosphorylation (PubMed:[30916345](#), PubMed:[31465518](#)). The nuclear import process is regulated by the small GTPase Ran that partitions between cytoplasm and nucleus in the predominantly GDP- and GTP-bound form, respectively (PubMed:[23878195](#), PubMed:[24449914](#)). Importin associates with target cargo proteins in the cytoplasm, and the competitive binding of GTP-bound Ran induces the release of cargos in the nucleus (PubMed:[23878195](#), PubMed:[24449914](#)).

Cellular Location

Nucleus envelope. Cytoplasm. Note=Localizes to the nuclear envelope and annulate lamellae, which consists in stacks of endoplasmic reticulum membranes containing a high density of nuclear pores

Tissue Location

Expressed in skeletal muscle.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.