

LD78 beta/CCL3L1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56989

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	P16619
Reactivity	Recombinant Fragment
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10161
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LD78 beta/CCL3L1
Epitope Specificity	31-93/93
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the intercrine beta (chemokine CC) family.
Post-translational modifications	The N-terminal processed forms LD78-beta(3-70) and LD78-beta(5-70) are produced by proteolytic cleavage after secretion from peripheral blood monocytes. The cleavage to yield LD78-beta(3-70) is probably achieved by DPP4.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene is one of several cytokine genes that are clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins that function in inflammatory and immunoregulatory processes. The protein encoded by this gene binds to several chemokine receptors, including chemokine binding protein 2 and chemokine (C-C motif) receptor 5 (CCR5). CCR5 is a co-receptor for HIV, and binding of this protein to CCR5 inhibits HIV entry. The copy number of this gene varies among individuals, where most individuals have one to six copies, and a minority of individuals have zero or more than six copies. There are conflicting reports about copy number variation of this gene and its correlation to disease susceptibility.[provided by RefSeq, Apr 2014]

Additional Information

Gene ID	414062;6349
Other Names	C-C motif chemokine 3-like 1, G0/G1 switch regulatory protein 19-2, LD78-beta(1-70), PAT 464.2, Small-inducible cytokine A3-like 1, Tonsillar lymphocyte LD78 beta protein, LD78-beta(3-70), LD78-beta(5-70), CCL3L1, D17S1718, G0S19-2, SCYA3L1
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500

Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CCL3L1
Synonyms	D17S1718, G0S19-2, SCYA3L1
Function	Chemotactic for lymphocytes and monocytes. Is a ligand for CCR1, CCR3 and CCR5. Is an inhibitor of HIV-1-infection. The processed form LD78-beta(3-70) shows a 20-fold to 30-fold higher chemotactic activity and is a very potent inhibitor of HIV-1-infection. LD78- beta(3-70) is also a ligand for CCR1, CCR3 and CCR5.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.