

LDLRAD3 Rabbit pAb

LDLRAD3 Rabbit pAb

Catalog # AP56990

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q86YD5
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37419
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LDLRAD3
Epitope Specificity	101-200/345
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane.
SIMILARITY	Belongs to the LDLR family. Contains 3 LDL-receptor class A domains.
SUBUNIT	Interacts with APP precursor C-terminus
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	143458
Other Names	Low-density lipoprotein receptor class A domain-containing protein 3, LDLR class A domain-containing protein 3, LDLRAD3 (HGNC:27046), LRAD3
Target/Specificity	Expressed at high levels in brain, lung, skeletal muscle, and pancreas. Expressed at moderate levels in heart, placenta, and kidney but not detected in the liver.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LDLRAD3 (HGNC:27046)
-------------	--

Synonyms

LRAD3

Function


May influence APP processing, resulting in a decrease in sAPP-alpha production and increased amyloidogenic P3 peptide production. May regulate ITCH and NEDD4 E3 ligase activity and degradation (PubMed:[26854353](https://pubmed.ncbi.nlm.nih.gov/26854353/)).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed at high levels in brain, lung, skeletal muscle, and pancreas.
Expressed at moderate levels in heart, placenta, and kidney but not detected in the liver

Images

Tissue/cell: human lung carcinoma; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal
goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-LDLRAD3 Polyclonal Antibody,
Unconjugated(AP56990) 1:500, overnight at 4°C, followed
by conjugation to the secondary antibody(SP-0023) and
DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.