

LINGO3 Rabbit pAb

LINGO3 Rabbit pAb

Catalog # AP57019

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P0C6S8
Predicted	Human, Mouse, Rat, Dog, Pig, Horse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64881
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LINGO3
Epitope Specificity	351-450/592
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane.
SIMILARITY	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 11 LRR (leucine-rich) repeats. Contains 1 LRRCT domain. Contains 1 LRRNT domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Lingo-3 is a 592 amino acid single-pass type I membrane protein. Lingo-3 contains one immunoglobulin-like C2-type domain and fourteen leucine-rich repeats and is glycosylated at various positions during post-translational modification. Lingo-3 is expressed in a broad but specific pattern in many tissues across the embryo. The gene encoding Lingo-3 maps to to human chromosome 19p13.3 and mouse chromosome 10 C1. Chromosome 19 consists of approximately 63 million bases and makes up over 2% of human genomic DNA. Chromosome 19 is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc alpha receptors.

Additional Information

Gene ID	645191
Other Names	Leucine-rich repeat and immunoglobulin-like domain-containing nogo receptor-interacting protein 3, Leucine-rich repeat neuronal protein 2, Leucine-rich repeat neuronal protein 6B, LINGO3, LERN2, LRRN6B
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	LINGO3
Synonyms	LERN2, LRRN6B
Cellular Location	Membrane; Single-pass type I membrane protein

Background

Lingo-3 is a 592 amino acid single-pass type I membrane protein. Lingo-3 contains one immunoglobulin-like C2-type domain and fourteen leucine-rich repeats and is glycosylated at various positions during post-translational modification. Lingo-3 is expressed in a broad but specific pattern in many tissues across the embryo. The gene encoding Lingo-3 maps to human chromosome 19p13.3 and mouse chromosome 10 C1. Chromosome 19 consists of approximately 63 million bases and makes up over 2% of human genomic DNA. Chromosome 19 is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin superfamily members including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc alpha receptors.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.