

LYRM4 Rabbit pAb

LYRM4 Rabbit pAb
Catalog # AP57164

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9HD34
Predicted	Rat, Human, Mouse, Rabbit, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10758
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LYRM4
Epitope Specificity	1-60/91
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion. Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	57128
Other Names	LYR motif-containing protein 4, LYRM4 (HGNC:21365), C6orf149, ISD11
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LYRM4 (HGNC:21365)
Synonyms	C6orf149, ISD11
Function	Stabilizing factor, of the core iron-sulfur cluster (ISC) assembly complex, that regulates, in association with NDUFB1, the stability and the cysteine desulfurase activity of NFS1 and participates in the [2Fe-2S] clusters assembly on the scaffolding protein ISCU (PubMed: 17331979 , PubMed: 31664822). The core iron-sulfur cluster (ISC) assembly complex is involved in the de novo

synthesis of a [2Fe-2S] cluster, the first step of the mitochondrial iron-sulfur protein biogenesis. This process is initiated by the cysteine desulfurase complex (NFS1:LYRM4:NDUFAB1) that produces persulfide which is delivered on the scaffold protein ISCU in a FXN-dependent manner. Then this complex is stabilized by FDX2 which provides reducing equivalents to accomplish the [2Fe-2S] cluster assembly. Finally, the [2Fe-2S] cluster is transferred from ISCU to chaperone proteins, including HSCB, HSPA9 and GLRX5 (By similarity). May also participates in the iron-sulfur protein biogenesis in the cytoplasm through its interaction with the cytoplasmic form of NFS1 (PubMed:[19454487](#)).

Cellular Location	Mitochondrion. Nucleus
Tissue Location	Reduced mRNA levels in Friedreich ataxia patients.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.