

MAFG Rabbit pAb

MAFG Rabbit pAb
Catalog # AP57180

Product Information

Application	WB
Primary Accession	O15525
Reactivity	Human
Predicted	Rat, Pig, Dog, Mouse, Chicken, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17850
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MAFG
Epitope Specificity	1-100/162
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Globin gene expression is regulated through nuclear factor erythroid-2 (NFE2) elements located in enhancer-like locus control regions positioned many kb upstream of alpha- and beta-gene clusters (summarized by Blank et al., 1997 [PubMed 9166829]). NFE2 DNA-binding activity consists of a heterodimer containing a ubiquitous small Maf protein (MafF, MIM 604877; MafG; or MafK, MIM 600197) and the tissue-restricted protein p45 NFE2 (MIM 601490). Both subunits are members of the activator protein-1-like superfamily of basic leucine zipper (bZIP) proteins (see MIM 165160).[supplied by OMIM, Mar 2010]

Additional Information

Gene ID	4097
Other Names	Transcription factor MafG, V-maf musculoaponeurotic fibrosarcoma oncogene homolog G, hMAF, MAFG
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	MAFG
Function	Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves (PubMed: 11154691). However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins, such as NFE2, NFE2L1 and NFE2L2, and recruiting them to specific DNA-binding sites (PubMed: 11154691 , PubMed: 8932385 , PubMed: 9421508). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NFE2L2 transcription factor (PubMed: 11154691). Transcription factor, component of erythroid- specific transcription factor NFE2L2 (PubMed: 11154691). Activates globin gene expression when associated with NFE2L2 (PubMed: 11154691). May be involved in signal transduction of extracellular H(+) (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00978, ECO:0000269 PubMed:11154691}
Tissue Location	Highly expressed in skeletal muscle. Also expressed in heart and brain

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Hela(Human) Cell Lysate at 40 ug
Jurkat(Human) Cell Lysate at 40 ug
Primary: Anti-MAFG (AP57180) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 18 kD
Observed band size: 18 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.