

MAK16/RBM13 Rabbit pAb

MAK16/RBM13 Rabbit pAb

Catalog # AP57192

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9BXY0
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35369
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MAK16/RBM13
Epitope Specificity	1-100/300
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear; nucleolus
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	RBM13 is a nucleolar protein containing 300 amino acids that belongs to the MAK16 family. The gene encoding RBM13 maps to human chromosome 8, which consists of nearly 146 million base pairs, encodes over 800 genes and is associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder, Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some leukemias and lymphomas are thought to occur as a result of defects in specific genes that map to chromosome 8.

Additional Information

Gene ID	84549
Other Names	Protein MAK16 homolog, NNP78, Protein RBM13, MAK16, RBM13
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	MAK16
-------------	-------

Synonyms	RBM13
Cellular Location	Nucleus, nucleolus

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MAK16/RBM13) Polyclonal Antibody, Unconjugated (AP57192) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.