

MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57208

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q9NS73</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 39281
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human MBIP/MAP3K12

binding inhibitory protein 1

Epitope Specificity 151-250/344

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus. Cytoplasm. Shows a cytoplasmic localization when coexpressed with

MAP3K12.

SUBUNIT Component of the ADA2A-containing complex (ATAC), composed of CSRP2BP,

KAT2A, TADA2L, TADA3L, ZZ3, MBIP, WDR5, YEATS2, CCDC101 and DR1. In the

complex, it probably interacts directly with KAT2A, CSRP2BP and WDR5.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Additional Information

Gene ID 51562

Other Names MAP3K12-binding inhibitory protein 1, MAPK upstream kinase-binding

inhibitory protein, MUK-binding inhibitory protein, MBIP

Target/Specificity Ubiquitous. High expression seen in the heart and lung.

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name MBIP

Function Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK

pathway. Component of the ATAC complex, a complex with histone

acetyltransferase activity on histones H3 and H4.

Cellular Location Nucleus. Cytoplasm. Note=Shows a cytoplasmic localization when

coexpressed with MAP3K12

Tissue Location Ubiquitous. High expression seen in the heart and lung

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.