

MASS1/GPR98 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57215

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q8WVG9
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	693069
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MASS1/GPR98
Epitope Specificity	2451-2550/6306
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane.
SIMILARITY	Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily. Contains 35 Calx-beta domains. Contains 6 EAR repeats. Contains 1 GPS domain.
DISEASE	Defects in GPR98 are the cause of Usher syndrome type 2C (USH2C) [MIM:605472]. USH is a genetically heterogeneous condition characterized by the association of retinitis pigmentosa with sensorineural deafness. Age at onset and differences in auditory and vestibular function distinguish Usher syndrome type 1 (USH1), Usher syndrome type 2 (USH2) and Usher syndrome type 3 (USH3). USH2 is characterized by congenital mild hearing impairment with normal vestibular responses. Defects in GPR98 may be a cause of familial febrile convulsions type 4 (FEB4) [MIM:604352]; also known as familial febrile seizures 4. Febrile convulsions are seizures associated with febrile episodes in childhood without any evidence of intracranial infection or defined pathologic or traumatic cause. It is a common condition, affecting 2-5% of children aged 3 months to 5 years. The majority are simple febrile seizures (generally defined as generalized onset, single seizures with a duration of less than 30 minutes). Complex febrile seizures are characterized by focal onset, duration greater than 30 minutes, and/or more than one seizure in a 24 hour period. The likelihood of developing epilepsy following simple febrile seizures is low. Complex febrile seizures are associated with a moderately increased incidence of epilepsy.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	MASS1 (for monogenic audiogenic seizure susceptibility 1) is one of the largest known GPCRs and is therefore referred to as Very Large G protein-coupled receptor 1 (VLGR1) (1,2). MASS1 is a large, calcium-binding GPCR expressed in the central nervous system and the eye (2,3). MASS1 has a large ectodomain containing multiple calcium exchanger beta repeats that resemble regulatory domains of sodium-calcium exchanger proteins (3). The human MASS1 gene maps to chromosome 5q14 and encodes a 1967 amino

acid protein (1,2,4). The MASS1 gene has been linked to the autosomal recessive inheritance of general epilepsy in Frings mice that have seizures in response to loud noises (5).

Additional Information

Gene ID	84059
Other Names	Adhesion G-protein coupled receptor V1, ADGRV1, 3.4.-., ADGRV1 (HGNC:17416)
Target/Specificity	Expressed at low levels in adult tissues.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ADGRV1 (HGNC:17416)
Function	G-protein coupled receptor which has an essential role in the development of hearing and vision. Couples to G-alpha(i)-proteins, GNAI1/2/3, G-alpha(q)-proteins, GNAQ, as well as G-alpha(s)-proteins, GNAS, inhibiting adenylate cyclase (AC) activity and cAMP production. Required for the hair bundle ankle formation, which connects growing stereocilia in developing cochlear hair cells of the inner ear. In response to extracellular calcium, activates kinases PKA and PKC to regulate myelination by inhibiting the ubiquitination of MAG, thus enhancing the stability of this protein in myelin-forming cells of the auditory pathway. In retina photoreceptors, the USH2 complex is required for the maintenance of periciliary membrane complex that seems to play a role in regulating intracellular protein transport. Involved in the regulation of bone metabolism.
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:Q8VHN7}; Multi-pass membrane protein {ECO:0000250 UniProtKB:Q8VHN7}. Cell projection, stereocilium membrane {ECO:0000250 UniProtKB:Q8VHN7} Photoreceptor inner segment {ECO:0000250 UniProtKB:Q8VHN7} Note=Localizes at the ankle region of the stereocilia. In photoreceptors, localizes at a plasma membrane microdomain in the apical inner segment that surrounds the connecting cilia called periciliary membrane complex. {ECO:0000250 UniProtKB:Q8VHN7}
Tissue Location	Expressed at low levels in adult tissues.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.