

# MBNL2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57221

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q5VZF2</a>
<b>Reactivity</b>	Rat, Dog, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	40518
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human MBNL2
<b>Epitope Specificity</b>	101-200/373
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Nucleus. Cytoplasm. Greater concentration in the nucleus. Expressed in or near large cytoplasmic adhesion plaques. Location in the cytoplasm is microtubule-dependent. In both DM1 and DM2 patients, colocalizes with nuclear foci of retained expanded-repeat transcripts.
<b>SIMILARITY</b>	Belongs to the muscleblind family. Contains 4 C3H1-type zinc fingers.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	This gene is a member of the muscleblind protein family which was initially described in <i>Drosophila melanogaster</i> . This gene encodes a C3H-type zinc finger protein that modulates alternative splicing of pre-mRNAs. Muscleblind proteins bind specifically to expanded dsCUG RNA but not to normal size CUG repeats and may thereby play a role in the pathophysiology of myotonic dystrophy. Several alternatively spliced transcript variants have been described but the full-length nature of only some have been determined. [provided by RefSeq, Mar 2012]

## Additional Information

---

<b>Gene ID</b>	10150
<b>Other Names</b>	Muscleblind-like protein 2, Muscleblind-like protein 1, Muscleblind-like protein-like, Muscleblind-like protein-like 39, MBNL2, MBLL, MBLL39, MLP1
<b>Target/Specificity</b>	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000

<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	MBNL2
<b>Synonyms</b>	MBLL, MBLL39, MLP1
<b>Function</b>	Mediates pre-mRNA alternative splicing regulation. Acts either as activator or repressor of splicing on specific pre-mRNA targets. Inhibits cardiac troponin-T (TNNT2) pre-mRNA exon inclusion but induces insulin receptor (IR) pre-mRNA exon inclusion in muscle. Antagonizes the alternative splicing activity pattern of CELF proteins. RNA-binding protein that binds to 5'ACACCC-3' core sequence, termed zipcode, within the 3'UTR of ITGA3. Binds to CUG triplet repeat expansion in myotonic dystrophy muscle cells by sequestering the target RNAs. Together with RNA binding proteins RBPMS and RBFOX2, activates vascular smooth muscle cells alternative splicing events (By similarity). Regulates NCOR2 alternative splicing (By similarity). Seems to regulate expression and localization of ITGA3 by transporting it from the nucleus to cytoplasm at adhesion plaques. May play a role in myotonic dystrophy pathophysiology (DM).
<b>Cellular Location</b>	Nucleus. Cytoplasm. Note=Greater concentration in the nucleus. Expressed in or near large cytoplasmic adhesion plaques (PubMed:16273094). Location in the cytoplasm is microtubule-dependent (PubMed:16273094). In both DM1 and DM2 patients, colocalizes with nuclear foci of retained expanded-repeat transcripts (PubMed:11929853)
<b>Tissue Location</b>	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.