

MIA3 Rabbit pAb

MIA3 Rabbit pAb
Catalog # AP57315

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q5JRA6
Reactivity	Rat, Mouse
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	213702
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MIA3
Epitope Specificity	1251-1350/1907
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum membrane. Localizes at endoplasmic reticulum exit sites. After loading of COL7A1 into transport carriers, it is not incorporated into COPII carriers and remains in the endoplasmic reticulum membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	375056
Other Names	Transport and Golgi organization protein 1 homolog, TANGO1, C219-reactive peptide, D320, Melanoma inhibitory activity protein 3 {ECO:0000312 HGNC:HGNC:24008}, MIA3 (HGNC:24008)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MIA3 (HGNC:24008)
Function	Plays a role in the transport of cargos that are too large to fit into COPII-coated vesicles and require specific mechanisms to be incorporated into membrane-bound carriers and exported from the endoplasmic

reticulum. This protein is required for collagen VII (COL7A1) secretion by loading COL7A1 into transport carriers. It may participate in cargo loading of COL7A1 at endoplasmic reticulum exit sites by binding to COPII coat subunits Sec23/24 and guiding SH3-bound COL7A1 into a growing carrier. Does not play a role in global protein secretion and is apparently specific to COL7A1 cargo loading. However, it may participate in secretion of other proteins in cells that do not secrete COL7A1. It is also specifically required for the secretion of lipoproteins by participating in their export from the endoplasmic reticulum (PubMed:[19269366](#), PubMed:[27138255](#)). Required for correct assembly of COPII coat components at endoplasmic reticulum exit sites (ERES) and for the localization of SEC16A and membrane-bound ER- resident complexes consisting of MIA2 and PREB/SEC12 to ERES (PubMed:[28442536](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Note=Localizes at endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:32101163). SEC16A is required for its proper localization to ERES. After loading of COL7A1 into transport carriers, it is not incorporated into COPII carriers and remains in the endoplasmic reticulum membrane.


Tissue Location

Broadly expressed, except in bone marrow and peripheral blood mononuclear cells. Down-regulated in melanoma tissue

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MIA3) Polyclonal Antibody, Unconjugated (AP57315) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.