

NCRNA00114 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57374

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q6XXX2 Human Reactivity Host Rabbit Clonality Polyclonal **Calculated MW** 15753 **Physical State** Liquid

KLH conjugated synthetic peptide derived from human NCRNA00114 **Immunogen**

Epitope Specificity 1-100/140 Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. **Important Note**

This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions The smallest of the human chromosomes, 21 makes up about 1.5% of the

> human genome. Chromosome 21 contains nearly 300 genes and 47 million base pairs. Down syndrome, also known as trisomy 21, is the disease most commonly associated with chromosome 21. Alzheimer's disease, Jervell and Lange-Nielsen syndrome and amyotrophic lateral sclerosis are also associated

with chromosome 21. Translocations are found to occur between

chromosome 21 and 8, and chromosome 21 and 12, in certain leukemias. The C21orf24 gene product has been provisionally designated C21orf24 pending

further characterization.

Additional Information

Other Names Putative uncharacterized protein encoded by LINC00114, LINC00114,

C21orf24, NCRNA00114

Target/Specificity Expressed in a range of cell lines, including B-cell lymphoma and prostate.

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce **Format**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When **Storage**

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name LINC00114


Synonyms C21orf24, NCRNA00114

Tissue Location Expressed in a range of cell lines, including B- cell lymphoma and prostate.

Images



Paraformaldehyde-fixed, paraffin embedded (human liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NCRNA00114) Polyclonal Antibody, Unconjugated (AP57374) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NCRNA00114) Polyclonal Antibody, Unconjugated (AP57374) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.