

MYO1D Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57421

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O94832
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116202

Additional Information

Gene ID	4642
Other Names	Unconventional myosin-Id, MYO1D, KIAA0727
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MYO1D
Synonyms	KIAA0727
Function	Unconventional myosin that functions as actin-based motor protein with ATPase activity (By similarity). Plays a role in endosomal protein trafficking, and especially in the transfer of cargo proteins from early to recycling endosomes (By similarity). Required for normal planar cell polarity in ciliated tracheal cells, for normal rotational polarity of cilia, and for coordinated, unidirectional ciliary movement in the trachea. Required for normal, polarized cilia organization in brain ependymal epithelial cells (By similarity).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q63357}. Perikaryon {ECO:0000250 UniProtKB:Q63357}. Cell projection, dendrite {ECO:0000250 UniProtKB:Q63357}. Early endosome {ECO:0000250 UniProtKB:F1PRN2}. Cytoplasm, cell cortex {ECO:0000250 UniProtKB:Q63357}. Note=Colocalizes with the actin cytoskeleton in the cell cortex close to the apical cell membrane Colocalizes

with cytoplasmic puncta that are reminiscent of transport vesicles.
{ECO:0000250|UniProtKB:Q63357}

Tissue Location

Expressed in many tissues. Highest levels in brain, followed by lung and ovary; expression is lowest in spleen

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.