10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

RNF208 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5743c

Product Information

Application	IHC-P, FC, WB, E
Primary Accession	Q9H0X6
Other Accession	NP_112587.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB27242
Calculated MW	27964
Antigen Region	86-115

Additional Information

Gene ID	727800
Other Names	RING finger protein 208, RNF208
Target/Specificity	This RNF208 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 86-115 amino acids from the Central region of human RNF208.
Dilution	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RNF208 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	RNF208

References

Simpson, J.C., et al. EMBO Rep. 1(3):287-292(2000)


Images



Western blot analysis of lysate from A431 cell line, using RNF208 Antibody (Center)(Cat. #AP5743c). AP5743c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.



RNF208 Antibody (Center) (Cat. #AP5743c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RNF208 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



RNF208 Antibody (Center) (Cat. #AP5743c) flow cytometric analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.