

NLRC5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57462

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q86WI3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	204595
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NLRC5
Epitope Specificity	651-750/1866
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
SIMILARITY	Belongs to the NLRP family. Contains 26 LRR (leucine-rich) repeats. Contains 1 NACHT domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the caspase recruitment domain-containing NLR family. This gene plays a role in cytokine response and antiviral immunity through its inhibition of NF-kappa-B activation and negative regulation of type I interferon signaling pathways. [provided by RefSeq, Oct 2011]


Additional Information

Gene ID	84166
Other Names	Protein NLRC5, Caterpillar protein 16.1, CLR16.1, Nucleotide-binding oligomerization domain protein 27, Nucleotide-binding oligomerization domain protein 4, NLRC5, NOD27, NOD4
Target/Specificity	Expressed in spleen, thymus, lung, brain, tonsil, heart and prostate.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NLRC5
Synonyms	NOD27, NOD4
Function	Probable regulator of the NF-kappa-B and type I interferon signaling pathways. May also regulate the type II interferon signaling pathway. Plays a role in homeostatic control of innate immunity and in antiviral defense mechanisms.
Cellular Location	Cytoplasm
Tissue Location	Expressed in spleen, thymus, lung, brain, tonsil, heart and prostate.

Images



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NLRC5) Polyclonal Antibody, Unconjugated (AP57462) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.