

# NOL4 Rabbit pAb

NOL4 Rabbit pAb  
Catalog # AP57471

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF
<b>Primary Accession</b>	<a href="#">O94818</a>
<b>Reactivity</b>	Mouse
<b>Predicted</b>	Human, Rat, Dog, Rabbit, Sheep
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	71357
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human NOL4
<b>Epitope Specificity</b>	1-100/638
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Nucleus > nucleolus.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	NOL4 is a 524 amino acid protein that is predominantly expressed in testis and brain. NOL4 contains at least two domains that direct it to subnuclear locations. The gene encoding NOL4 is located on human chromosome 18, which houses over 300 protein-coding genes and contains nearly 76 million bases. There are a variety of diseases associated with defects in chromosome 18-localized genes, some of which include Trisomy 18 (also known as Edwards syndrome), Niemann-Pick disease, hereditary hemorrhagic telangiectasia, erythropoietic protoporphyrin and follicular lymphomas. There are two isoforms of NOL4 that are produced as a result of alternative splicing events.

## Additional Information

---

<b>Gene ID</b>	8715
<b>Other Names</b>	Nucleolar protein 4, Nucleolar-localized protein, NOL4, NOLP
<b>Target/Specificity</b>	Expressed predominantly in fetal brain, adult brain and testis.
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	NOL4
<b>Synonyms</b>	NOLP
<b>Cellular Location</b>	Nucleus, nucleolus
<b>Tissue Location</b>	Expressed predominantly in fetal brain, adult brain and testis.


## Background

---

NOL4 is a 524 amino acid protein that is predominantly expressed in testis and brain. NOL4 contains at least two domains that direct it to subnuclear locations. The gene encoding NOL4 is located on human chromosome 18, which houses over 300 protein-coding genes and contains nearly 76 million bases. There are a variety of diseases associated with defects in chromosome 18-localized genes, some of which include Trisomy 18 (also known as Edwards syndrome), Niemann-Pick disease, hereditary hemorrhagic telangiectasia, erythropoietic protoporphyrin and follicular lymphomas. There are two isoforms of NOL4 that are produced as a result of alternative splicing events.

## Images

---



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NOL4) Polyclonal Antibody, Unconjugated (AP57471) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.