

# NSUN5P2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57541

## Product Information

---

|                          |                          |
|--------------------------|--------------------------|
| <b>Application</b>       | IHC-P, IHC-F, IF, ICC, E |
| <b>Primary Accession</b> | <a href="#">Q63ZY6</a>   |
| <b>Reactivity</b>        | Human                    |
| <b>Host</b>              | Rabbit                   |
| <b>Clonality</b>         | Polyclonal               |
| <b>Calculated MW</b>     | 34347                    |

## Additional Information

---

|                    |                                                                                                                                                                                                 |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Other Names</b> | Putative methyltransferase NSUN5C, 2.1.1.-, NOL1/NOP2/Sun domain family member 5C, Williams-Beuren syndrome chromosomal region 20C protein, NSUN5P2, NSUN5C, WBSCR20B, WBSCR20C                 |
| <b>Dilution</b>    | IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000                                                                                                                   |
| <b>Storage</b>     | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

## Protein Information

---

|                        |                                                                         |
|------------------------|-------------------------------------------------------------------------|
| <b>Name</b>            | NSUN5P2                                                                 |
| <b>Synonyms</b>        | NSUN5C, WBSCR20B, WBSCR20C                                              |
| <b>Function</b>        | May have S-adenosyl-L-methionine-dependent methyl-transferase activity. |
| <b>Tissue Location</b> | Ubiquitous.                                                             |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.