

TCF4 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5756a

Product Information

Application	WB, FC, E
Primary Accession	P15884
Other Accession	Q60722 , NP_003190.1
Reactivity	Human, Rat, Mouse
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB21346
Calculated MW	71308
Antigen Region	48-73

Additional Information

Gene ID	6925
Other Names	Transcription factor 4, TCF-4, Class B basic helix-loop-helix protein 19, bHLHb19, Immunoglobulin transcription factor 2, ITF-2, SL3-3 enhancer factor 2, SEF-2, TCF4, BHLHB19, ITF2, SEF2
Target/Specificity	This TCF4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 48-73 amino acids from the N-terminal region of human TCF4.
Dilution	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TCF4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TCF4
Synonyms	BHLHB19, ITF2, SEF2

Function	Transcription factor that binds to the immunoglobulin enhancer Mu-E5/KE5-motif. Involved in the initiation of neuronal differentiation. Activates transcription by binding to the E box (5'- CANNTG-3'). Binds to the E-box present in the somatostatin receptor 2 initiator element (SSTR2-INR) to activate transcription (By similarity). Preferentially binds to either 5'-ACANNTGT-3' or 5'- CCANNTGG-3'.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00981, ECO:0000269 PubMed:22777675}
Tissue Location	Expressed in adult heart, brain, placenta, skeletal muscle and to a lesser extent in the lung. In developing embryonic tissues, expression mostly occurs in the brain


Background

TCF4 is transcription factor 4, a basic helix-turn-helix transcription factor. The encoded protein recognizes an Ephrussi-box ('E-box') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is expressed predominantly in pre-B-cells, although it is found in other tissues as well. Defects in this gene are a cause of Pitt-Hopkins syndrome.


References

Rosenfeld, J.A., et al. Genet. Med. 11(11):797-805(2009)
Herbst, A., et al. Biochem. Biophys. Res. Commun. 387(4):736-740(2009)
Rena, V., et al. Placenta 30(10):876-883(2009)
Henthorn, P., et al. Science 247(4941):467-470(1990)

Images



TCF4 Antibody (N-term) (Cat. #AP5756a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the TCF4 antibody detected the TCF4 protein (arrow).



TCF4 Antibody (N-term) (Cat. #AP5756a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.