

NUP155 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57574

Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O75694
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	155199

Additional Information

Gene ID	9631
Other Names	Nuclear pore complex protein Nup155, 155 kDa nucleoporin, Nucleoporin Nup155, NUP155, KIAA0791
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	NUP155
Synonyms	KIAA0791
Function	Essential component of nuclear pore complex. Could be essential for embryogenesis. Nucleoporins may be involved both in binding and translocating proteins during nucleocytoplasmic transport.
Cellular Location	Nucleus, nuclear pore complex {ECO:0000250 UniProtKB:P37199}. Nucleus membrane {ECO:0000250 UniProtKB:P37199}; Peripheral membrane protein {ECO:0000250 UniProtKB:P37199}; Cytoplasmic side {ECO:0000250 UniProtKB:P37199}. Nucleus membrane {ECO:0000250 UniProtKB:P37199}; Peripheral membrane protein {ECO:0000250 UniProtKB:P37199}; Nucleoplasmic side {ECO:0000250 UniProtKB:P37199}. Note=In mitosis, assumes a diffuse cytoplasmic distribution probably as a monomer, before reversing back into a punctate nuclear surface localization at the end of mitosis {ECO:0000250 UniProtKB:P37199}

Tissue Location

Expressed in all tissues tested, including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NUP155) Polyclonal Antibody, Unconjugated (AP57574) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.